

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Romaguera, D; Guevara, M; Norat, T; Langenberg, C; Forouhi, NG; Sharp, S; Slimani, N; Schulze, MB; Buijsse, B; Buckland, G; Molina-Montes, E; Sanchez, MJ; Moreno-Iribas, MC; Bendinelli, B; Grioni, S; van der Schouw, YT; Arriola, L; Beulens, JW; Boeing, H; Clavel-Chapelon, F; Cottet, V; Crowe, FL; de Lauzon-Guillan, B; Franks, PW; Gonzalez, C; Hallmans, G; Kaaks, R; Key, TJ; Khaw, K; Nilsson, P; Overvad, K; Palla, L; Palli, D; Panico, S; Quiros, JR; Rolandsson, O; Romieu, I; Sacerdote, C; Spijkerman, AMW; Teucher, B; Tjonneland, A; Tormo, MJ; Tumino, R; van der, ADL; Feskens, EJM; Riboli, E; Wareham, NJ; Interact, C (2011) Mediterranean diet and type 2 diabetes risk in the European Prospective Investigation into Cancer and Nutrition (EPIC) study: the InterAct project. *Diabetes care*, 34 (9). pp. 1913-8. ISSN 0149-5992

Downloaded from: <http://researchonline.lshtm.ac.uk/1924530/>

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: Creative Commons Attribution Non-commercial No Derivatives
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

SUPPLEMENTARY DATA

Author list:

Romaguera D, PhD (1), Guevara M, MD (2), Norat T, PhD (1), Langenberg C, PhD (3), Forouhi NG, FFPH (3), Sharp S, MSc (3), Slimani N, PhD (4), Schulze MB, PhD (5), Buijsse B, PhD (6), Buckland G, PhD (7), Molina-Montes E, PhD (8,9), Sánchez MJ, PhD (8,9), Moreno-Iribas MC, MD (2), Bendinelli B, PhD (10), Grioni S, MSc (11), van der Schouw YT, PhD (12), Arriola L, MD (9,13,14), Beulens JW, PhD (12), Boeing H, Prof (6), Clavel-Chapelon F, PhD (15), Cottet V, PhD (16), Crowe FL, PhD (17), de Lauzon-Guillan B, PhD (15,18), Franks PW, PhD (19), Gonzalez C, PhD (7), Hallmans G, PhD (20), Kaaks R, Prof (21), Key TJ, Prof (17), Khaw K, Prof (22), Nilsson P, Prof (19), Overvad K, Prof (23), Palla L, PhD (3), Palli D, MD (10), Panico S, MD (24), Quirós JR, PhD (25), Rolandsson O, PhD (20), Romieu I, PhD (4), Sacerdote C, PhD (26,27), Spijkerman AMW, PhD (28), Teucher B, PhD (21), Tjonneland A, PhD (29), Tormo MJ, PhD, (30), Tumino R, MD (31), van der A DL, PhD (28), Feskens EJM, Prof (32), Riboli E, Prof (1), Wareham NJ, Prof (3).

Affiliations:

(1) Department of Epidemiology & Biostatistics, School of Public Health, Imperial College London, St Mary's Campus, Norfolk Place, London W2 1PG, United Kingdom

(2) Public Health Institute of Navarra, Leyre 15, 31003 Pamplona, Spain and Public Health (CIBER Epidemiología y Salud Pública - CIBERESP)

(3) MRC Epidemiology Unit, Institute of Metabolic Science, Addenbrooke's Hospital, Box 285, Cambridge, CB2 0QQ, United Kingdom

(4) International Agency for Research of Cancer, 150 Cours Albert Thomas, 69372 Cedex 08 Lyon, France

(5) Department of Molecular Epidemiology, German Institute of Human Nutrition Potsdam Rehbruecke, Arthur-Scheunert-Allee 114-116, 14558 Nuthetal, Germany

(6) Department of Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Arthur-Scheunert-Allee 114-116, 14558 Nuthetal, Germany

(7) Unit Nutrition, Environment and Cancer, Department of Epidemiology, Catalan Institute of Oncology (ICO-IDIBELL), Gran Via s/n km 2,7, L'Hospitalet de Llobregat, 08907 Barcelona, Spain

(8) Andalusian School of Public Health, Cuesta del Observatorio 4, Ap. Correos 2070, 18080 Granada, Spain

(9) Consortium for Biomedical Research in Epidemiology and Public Health (CIBER Epidemiología y Salud Pública - CIBERESP), Spain

(10) Molecular&Nutritional Epidemiology Unit, Cancer Research and Prevention Institute (ISPO), Via delle Oblate 2, 50141, Florence, Italy

(11) Fondazione IRCCS Istituto Nazionale Tumori Milan, Via Venezian, 1, 20133 Milan, Italy

(12) Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Stratenum 6.131, PO Box 85.500, 3508 GA Utrecht, The Netherlands

(13) Public Health Division of Gipuzkoa, Basque Government, Av. Navarra 4, 20013 San Sebastian, Spain

(14) Instituto BIO-Donostia, Basque Government, Spain

(15) Inserm, CESP Centre for research in Epidemiology and Population Health, U1018: Nutrition, hormones and women's health, IGR, 39 rue Camille Desmoulins, 94800 Villejuif cedex, France

(16) Inserm U866, team 6, Burgundy Registry of Digestive Cancers, 7 Bd Jeanne d'Arc, BP 87900, 21079 Dijon Cedex, France

(17) Cancer Epidemiology Unit, University of Oxford, Richard Doll Building, Nuffield Department of Clinical Medicine, Roosevelt Drive, Oxford OX3 7LF, United Kingdom

(18) University Paris Sud 11, UMR5 1018, 94807 Villejuif cedex, France

SUPPLEMENTARY DATA

- (19) Department of Clinical Sciences, Clinical Research Center, Skåne University Hospital - Malmö (UMAS), Lund University, 20502 Malmö, Sweden
- (20) Department of Public Health and Clinical Medicine, Umea University, 90187 Umea, Sweden
- (21) German Cancer Research Centre, Im Neuenheimer Feld 280, 69120 Heidelberg, Germany
- (22) Department of Public Health and Primary Care, University of Cambridge Addenbrooke's Hospital Box 251 Cambridge CB2 2QQ UK, United Kingdom
- (23) Department of Epidemiology, School of Public Health, Bartholins Allé 2, DK-8000 Aarhus C, Denmark
- (24) Dipartimento di Medicina Clinica e Sperimentale, Università Federico II, Department of Clinical and Experimental Medicine, Federico II University, via Pansini 5 - 80131 Napoli, Italy
- (25) Consejería de Salud y Servicios Sanitarios, C/Ciriaco Miguel Vigil 9, 33006- Oviedo-Asturias, Spain
- (26) Center for Cancer Prevention (CPO-Piemonte), Via Santena 7, 10126, Torino, Italy
- (27) Human Genetic Foundation (HuGeF), Unit of Epidemiology and Molecular Genetics Via Nizza 52, 10126 Turin, Italy
- (28) National Institute for Public Health and the Environment (RIVM), PO Box 1, 3720 BA Bilthoven, The Netherlands
- (29) Danish Cancer Society, Strandboulevarden 49, 2100 Copenhagen, Denmark
- (30) Servicio de Epidemiología, Consejería de Sanidad y Consumo, C/ Luis Fontes Pagán, nº9, 1ª Planta, (Edif. EMI Hospital Universitario "Reina Sofía"), 30003 Murcia, Spain
- (31) Cancer Registry and Histopathology Unit, "Civile - M.P. Arezzo" Hospital, ASP 7 Via Dante Nr. 109, 97100 Ragusa, Italy
- (32) Division of Human Nutrition - Section Nutrition and Epidemiology, University of Wageningen, PO Box 8129, 6700 EV Wageningen, The Netherlands

Authors' contributions:

NJW is the principal investigator of InterAct and is the guarantor of the paper. ER is the overall coordinator of the EPIC study, which was conceptualized, designed and implemented in collaboration with the main investigators in the collaborating countries. EJMF is the leader of the Nutrition and Physical Activity Research Line of InterAct. DR and TN designed this analysis. MG analysed the data. DR wrote the manuscript with close assistance from working team members: MG, TN, CL, NF, SS, NS, MS, BB, GB, EM-M, M-JS, MCM-I, BB, SG, YTvdS, EJMF, ER and NJW. All authors contributed to either EPIC or InterAct data collection, study management, or coordination. All authors contributed to data interpretation and critical reading of the manuscript and have seen and approved the final version of the manuscript.

SUPPLEMENTARY DATA

Supplementary Table 1. Distribution of cases and characteristics of the InterAct sample by country

		Subcohort		Non - subcohort		Total sample		
Country	N	Person- years	Cases (%)	Cases	N	Female, %	Age at enrolment, mean (SD)	rMED score, mean (SD)
France	581	5,838.49	9 (1.5)	271	852	100.0	56.9 (6.5)	9.2 (2.5)
Italy	1,952	23,521.82	58 (3.0)	1,287	3,239	64.9	51.3 (7.7)	11.1 (2.4)
Spain	3,513	45,359.78	254 (7.2)	2,240	5,753	56.5	50.4 (7.8)	11.0 (2.5)
UK-General population	1,064	12,231.67	28 (2.6)	826	1,890	51.0	60.0 (9.2)	8.2 (2.6)
UK-Health conscious	234	2,700.93	0 (0.0)	128	362	74.0	50.9 (12.1)	11.0 (2.2)
Netherlands	1,462	17,533.49	40 (2.7)	767	2,229	80.5	53.7 (10.2)	6.4 (2.3)
Germany	2,024	21,803.36	60 (3.0)	1,490	3,514	50.2	52.4 (8.3)	7.4 (2.3)
Sweden	2,887	36,962.52	160 (5.5)	2,385	5,272	52.4	54.9 (9.7)	6.2 (2.4)
Denmark	2,081	23,052.02	140 (6.7)	1,851	3,932	44.3	57.0 (4.4)	6.8 (2.6)
Total	15,798	189,004.08	749 (4.7)	11,245	27,043	57.3	53.8 (8.7)	8.4 (3.2)

Abbreviations: rMED: relative Mediterranean diet score. SD: Standard deviation

SUPPLEMENTARY DATA

Supplementary Table 2. Multiple adjusted hazard ratios of type 2 diabetes associated with two unit increment in the relative-Mediterranean Diet score (rMED) and after alternate subtraction of each of its components

Dietary variable	HR *	95%CI	P-value †
rMED overall	0.96	0.94 - 0.99	0.002
rMED minus vegetables	0.95	0.93 - 0.98	<0.001
rMED minus fruit and nuts	0.96	0.94 - 0.99	0.003
rMED minus legumes	0.96	0.94 - 0.99	0.002
rMED minus fish	0.95	0.93 - 0.98	<0.001
rMED minus cereals	0.95	0.93 - 0.98	<0.001
rMED minus olive oil	0.97	0.95 - 1.00	0.027
rMED minus alcohol	0.98	0.95 - 1.01	0.121
rMED minus meat & meat products	0.98	0.95 - 1.00	0.058
rMED minus dairy products	0.96	0.93 - 0.99	0.001

* Modified Cox proportional hazard regression models stratified by center, and adjusted for sex, body mass index (as a continuous variable), educational level (no formal education, primary school, technical/professional school, secondary school and longer education including university degree), physical activity (inactive, moderately inactive, moderately active and active), smoking status (never, former and three categories of current smoker: 1-10 cigarettes day⁻¹, 11-20 cigarettes day⁻¹ and >20 cigarettes day⁻¹), total calorie intake (as a continuous variable) and corresponding subtracted rMED components (codified like in the score). Originally estimated logarithms of type 2 diabetes ratios were multiplied by 17/19 and then exponentiated to correct for eighteen point scale (except for rMED overall).

† Calculated from the original numbers.

SUPPLEMENTARY DATA

Supplementary Table 3. Sensitivity analyses for risk type 2 diabetes associated with a two-point increment in the relative-Mediterranean Diet score (rMED)

	No. Cases / No. Sub-cohort *	HR †	95%CI
Further adjusting by			
WC ‡	11,158 / 14,089	0.97	0.94 – 0.99
Hypertension	11,994 / 15,049	0.95	0.93 – 0.98
Hyperlipidaemia §	9,449 / 12,322	0.95	0.93 – 0.98
WC, hypertension and hyperlipidaemia ‡ §	9,449 / 12,322	0.96	0.93 – 0.99
Excluding			
The first two years of follow-up	10,969 / 14,983	0.96	0.94 – 0.99
Participants with chronic diseases at baseline §	3,823 / 9,602	0.95	0.91 – 0.98
The first two years of follow-up and participants with chronic diseases at baseline §	3,565 / 9,568	0.94	0.91 – 0.98
Miss-reporters of energy according to the Goldberg classification	7,605 / 10,580	0.96	0.93 – 1.00
Additive calibration			
Calibrated rMED score	11,994 / 15,049	0.97	0.94 – 0.99

Abbreviations: WC: waist circumference.

* No. in the Sub-cohort excluding T2DM cases.

† Modified Cox proportional hazard regression models stratified by center and adjusted for sex, body mass index (as a continuous variable), educational level (no formal education, primary school, technical/professional school, secondary school and longer education including university degree), physical activity (inactive, moderately inactive, moderately active and active), smoking status (never, former and three categories of current smoker: 1-10 cigarettes day⁻¹, 11-20 cigarettes day⁻¹ and >20 cigarettes day⁻¹) and total calorie intake (as a continuous variable).

‡ Excludes Umea where WC was not recorded.

§ Excludes Malmo and Umea where hyperlipidaemia was not asked.

|| Participants who reported cardiovascular diseases (myocardial infarction, angina or stroke), hypertension, hyperlipidemia or were obese (BMI>30 kg.m⁻²) at baseline.


SUPPLEMENTARY DATA

Supplementary Figure 1. Country-specific hazard ratios of type 2 diabetes associated with a two-point increment in the relative-Mediterranean Diet score (rMED).

Hazard Ratios and 95% CI derived by modified Cox proportional hazard regression models stratified by center and adjusted sex, body mass index (as a continuous variable), educational level (no formal education, primary school, technical/professional school, secondary school and longer education including university degree), physical activity (inactive, moderately inactive, moderately active and active), smoking status (never, former and three categories of current smoker: 1-10 cigarettes day⁻¹, 11-20 cigarettes day⁻¹ and >20 cigarettes day⁻¹) and total calorie intake (as a continuous variable). Weights are from random effects analysis.

Abbreviations: UK-HC: UK Health conscious group; UK-GP: UK General population.

SUPPLEMENTARY DATA



SUPPLEMENTARY DATA

Supplementary Figure 2. Multivariable hazard ratio of type 2 diabetes associated with a two-point increment in the relative-Mediterranean Diet Score (rMED) by country, according to the mean age at recruitment in the sub-cohort.

Area of symbol proportional to sample size (% weight).

Abbreviations: UK-HC: UK Health conscious group; UK-GP: UK General population

